

10089889 0898203

JC13 Rec'd PCT/PTC 25 MAR 2002

Practitioner's Docket No. U 013943-5

CHAPTER II

IN THE UNITED STATES ELECTED OFFICE (EO/US)

PCT/IN00/00094 27 SEPTEMBER 2000 28 SEPTEMBER 1999  
INTERNATIONAL APPLICATION NO. INTERNATIONAL FILING DATE PRIORITY DATE CLAIMED

CONTROLLED RELEASE COMPOSITIONS COMPRISING NIMESULIDE

TITLE OF INVENTION

1. Amarjit SINGH; 2. Rajesh JAIN

APPLICANT(S)

Box PCT  
Assistant Commissioner for Patents  
Washington, D.C. 20231  
ATTENTION: EO/US

PRELIMINARY AMENDMENT

Please amend the above identified application as follows:

IN THE CLAIMS :

Please cancel claims 12 and 13.


Please amend claims 3 and 4 as follows:

CERTIFICATE UNDER 37 1.10

I hereby certify that this paper is being deposited with the United States Postal Service on this date MARCH 25, 2002 in an envelope as "EXPRESS MAIL POST OFFICE TO ADDRESSEE" Mailing Label Number EV011021240US addressed to the: Assistant Commissioner for Patents, Washington, D.C. 20231

CONNIE YANNOTTI

(Type or print name of person mailing paper)

  
(Signature of person mailing paper)

NOTE: Each paper or fee referred to as enclosed herein has the number of the "EXPRESS MAIL" mailing label placed thereon prior to mailing 37 CFR 1.16(b).

3. (Amended) A controlled release pharmaceutical composition of nimesulide as claimed in claim 1 wherein there is no loss of bioavailability in comparison to an immediate release composition.

4. (Amended) A controlled release pharmaceutical composition of nimesulide as claimed in claim 1 wherein the sustaining materials are selected from the group comprising cellulose and cellulose derivatives, waxes, carbomers, polyalkylene polyols, polycarbophils, methacrylic acid derivatives, gelatins, gums, polyethylene oxides.

Please add new claims 14-17 as follows:

14. (New) A controlled release pharmaceutical composition of nimesulide as claimed in claim 2 wherein there is no loss of bioavailability in comparison to an immediate release composition.

15. (New) A controlled release pharmaceutical composition of nimesulide as claimed in claim 2 wherein the sustaining materials are selected from the group comprising cellulose and cellulose derivatives, waxes, carbomers, polyalkylene polyols, polycarbophils, methacrylic acid derivatives, gelatins, gums, polyethylene oxides.

16. (New) A controlled release pharmaceutical composition of nimesulide as claimed in claim 3 wherein the sustaining materials are selected from the group comprising cellulose and cellulose derivatives, waxes, carbomers, polyalkylene polyols, polycarbophils, methacrylic acid derivatives, gelatins, gums, polyethylene oxides.

17. (New) A controlled release pharmaceutical composition of nimesulide as claimed in claim 14 wherein the sustaining materials are selected from the group comprising cellulose and cellulose derivatives, waxes, carbomers, polyalkylene polyols, polycarbophils, methacrylic acid derivatives, gelatins, gums, polyethylene oxides.

**Remarks**

The above amendatory action is taken solely for the purpose of avoiding claim fees that would otherwise accrue due to the presence of multiple dependent claims.

Respectfully submitted,



WILLIAM R. EVANS  
LADAS & PARRY  
26 WEST 61<sup>ST</sup> STREET  
NEW YORK, NEW YORK 10023  
REG.NO.25,858 (212)708-1930

**MARKED UP COPY**

3. (Amended) A controlled release pharmaceutical composition of nimesulide as claimed in claim 1 [and 2] wherein there is no loss of bioavailability in comparison to an immediate release composition.

4. (Amended) A controlled release pharmaceutical composition of nimesulide as claimed in claim 1 [to 3] wherein the sustaining materials are selected from the group comprising cellulose and cellulose derivatives, waxes, carbomers, polyalkylene polyols, polycarbophils, methacrylic acid derivatives, gelatins, gums, polyethylene oxides.

Controlled Release Compositions Comprising Nimesulide

The present invention relates to a controlled release composition of Nimesulide. The composition is related to a once-a-day dosage forms which are very useful in treatment of chronic diseases such as arthritis.

**TECHNICAL BACKGROUND OF THE INVENTION**

Nimesulide is a nonsteroidal anti-inflammatory drug (NSAID) that also has antipyretic and analgesic properties. The compound is weakly acidic (pKa = 6.5) and differs from other NSAIDs in that its chemical structure contains a sulfonanilide moiety as the acidic group. (fig. 1) (Magni E, Nimesulide an overview, Drug Invest 1991; 3 Suppl. 2: 1-3).




Fig. 1

The therapeutic effects of NSAIDs are largely the result of their ability to inhibit prostaglandin synthesis via inhibition of cyclo-oxygenase. Unfortunately, this effect is also responsible for the inhibition of gastroprotective prostaglandins, which leads to gastrointestinal intolerance.

In vitro, Nimesulide is a relatively weak inhibitor of prostaglandin synthesis and appears to exert its effects through a variety of mechanisms. (Magni E. The effect of nimesulide on prostanoid formation. *Drugs* 1993; 46 Suppl. 1:10-4.) Indeed, the mechanism of action of this drug is more complex than previously thought and may involve interference with the production/action of mediators other than prostaglandins such as enzymes, toxic oxygen derivatives, cytokines, platelet-activating factor (PAF) and histamine.

The anti-inflammatory, analgesic and antipyretic activities of Nimesulide, a non-steroidal anti-inflammatory drug (NSAID) of the sulfonanilide class, have been demonstrated in a number of experimental models and in numerous clinical trials. Nimesulide has exhibited potency similar to or greater than that of indomethacin, diclofenac, piroxicam and ibuprofen in standard animal models of inflammation such as carrageenin-induced rat paw oedema and inflammation, ultraviolet light-induced erythema in guinea-pigs and adjuvant arthritis in rats. The analgesic potency in nimesulide was similar to that of ibuprofen and less than that of indomethacin in an acetic acid writhing test in rats, and acetic acid and acetylcholine writhing tests in mice. Nimesulide has shown superior antipyretic potency to indomethacin, ibuprofen, aspirin and paracetamol (acetaminophen) in rats with yeast-induced fever.

Nimesulide is a relatively weak inhibitor of prostaglandin synthesis in vitro and appears to exert its effects through a variety of mechanisms including free-radical scavenging, effects on histamine release, the neutrophil myeloperoxidase pathway, bradykinin activity, tumour necrosis factor- $\alpha$

release, cartilage degradation, metalloprotease synthesis, phosphodiesterase type IV inhibition, platelet aggregation and synthesis of platelet activating factor. Animal studies have suggested that Nimesulide is less ulcerogenic than aspirin, indomethacin, naproxen, piroxicam and ibuprofen. Nimesulide appears to have little effect on renal prostaglandin synthesis in rats.

After oral administration of nimesulide 50 to 200 mg to healthy adult volunteers, peak serum concentrations of 1.98 to 9.85 mg/L are achieved within 1.22 to 3.17 hours. Compared with values obtained with oral drug administration, peak serum concentrations are slightly lower (2.14 to 2.32 mg/L) and are achieved more slowly (3 to 4.58 h) after rectal administration of nimesulide 100 and 200 mg. Oral drug absorption is nearly complete and concomitant administration of food may decrease the rate, but not the extent of absorption of nimesulide. The drug is extensively bound (99%) to plasma proteins and has an estimated apparent volume of distribution of 0.19 to 0.35 L/kg following oral administration.

In children, nimesulide suspension, granules and suppositories are more effective than placebo and are at least as effective as paracetamol, diclofenac, naproxen, lysine acetylsalicylate, mefenamic acid, ketoprofen and dipyrone in reducing in pain, inflammation and fever associated with respiratory tract infection, postoperative pain and musculoskeletal injury.

Nimesulide has been well tolerated by both young and elderly adults and children in 2 large postmarketing surveillance surveys. As with other NSAIDs, the most common adverse effects are gastrointestinal disturbances

(epigastralgia, heartburn, nausea, diarrhoea and vomitings 5.1 to 8.5% of patients), dermatological reactions (rash, pruritus; 0.2 to 0.6%) and central nervous system effects (dizziness, somnolence, headache; 0.3 to 0.4%). Withdrawal rates associated with short term (up to 30 days) nimesulide treatment range from 1.1 to 2.2% in adult, elderly and paediatric patients.

Available data indicate that the gastrointestinal tolerability of nimesulide in adults and children is similar to that of other NSAIDs. The rate of endoscopically verified gastroduodenal irritation with nimesulide appears to be similar to that with placebo and diclofenac and less than that with indomethacin. The drug is well tolerated by most patients intolerant of aspirin and/or other NSAIDs and by patients with asthma.

The literature surveys shows that different dosage forms reported for nimesulide are tablets, granules, suppositories and suspension (Drugs 48 (3): 431-454, 1994) and lately our group has patented transdermal (US Pat. No. 5688829) and intramuscular injection (US Pat. No. 5716609) formulations. The reported dosage forms have to be administered twice-a-day based on biological half life of nimesulide. The usual oral/rectal dosage of nimesulide in adults is 100 to 200 mg twice daily, orally. For treatment of chronic diseases like arthritis the twice daily dosing regimen is difficult to comply with.

One approach to improve the possible non-compliance with the regimen is to develop controlled release dosage form for nimesulide. The once-a-day dosage form is expected to significantly increase the dosing convenience and



patient compliance. However, controlled release once-a-day dosage form of nimesulide have not been reported so far.

Controlled release compositions for oral use in the form of matrix type monolithic tablets, beads, capsules and coated tablets are known. However poorly soluble drugs like nimesulide are known to give erratic and variable release under in-vivo conditions from such dosage forms.

One approach to formulate modified release dosage forms of NSAIDs is described in U.S. Pat. No. WO9912524, wherein a unit dosage form comprising two fractions (i) a first quick release fraction, and (ii) a second fraction containing coated delayed release multiple units is described. However, such dosage forms having different fractions and coated multiple units are difficult to prepare and very cost intensive. Moreover compression of such coated multiple units into tablets cause fracturing of the coat layer, thereby causing loss of reproducibility.

In U.S. Pat. No. 5788987 Buseti et.al. describe a time-specific controlled release dosage form. Such dosage forms are designed to provide delayed release of the active ingredient rather than extended release. Such formulations are not suitable for day long management of the disease.

## **SUMMARY OF THE INVENTION**

By expenditure of considerable intellectual effort and careful experimentation the inventors have discovered that nimesulide can be formulated into a controlled release once-a-day oral dosage form.

Such dosage forms provide extended release of nimesulide in-vivo when given once daily with reproducible bioavailability. Further the release of such dosage forms is not effected by pH changes in the gastrointestinal system.

The composition in accordance with present invention comprises a controlled release pharmaceutical composition of Nimesulide which comprises nimesulide as an active drug from 5% to 95% w/w of the composition in micronized form, one or more release sustaining materials from 2% to 95% w/w of the composition and pharmaceutical excipients from 0% to 90% w/w of the composition.

Preferably the composition in accordance with the present invention comprises nimesulide as an active drug from 20% to 70% w/w of the composition, one or more sustaining materials from 5% to 65% w/w of the composition and pharmaceutical excipients from 10% to 70% w/w of the composition.

More preferably the composition in accordance with the present invention comprises nimesulide as an active drug from 40% to 60% w/w of the composition, one or more sustaining materials from 8% to 20% w/w of the

composition and pharmaceutical excipients from 30% to 60% w/w of the composition.

## DETAILED DESCRIPTION OF INVENTION

In accordance with the present invention there is disclosed a controlled release composition of Nimesulide.

The composition in accordance with present invention comprises a controlled release pharmaceutical composition of Nimesulide which comprises nimesulide as an active drug from 5% to 95% w/w of the composition, one or more sustaining materials from 2% to 95% w/w of the composition and pharmaceutical excipients from 0% to 90% w/w of the composition. In another aspect, such compositions contain nimesulide in micronized form having average particle size below 5 microns.

Preferably the composition in accordance with the present invention comprises nimesulide as an active drug from 20% to 70% w/w of the composition, one or more sustaining materials from 5% to 65% w/w of the composition and pharmaceutical excipients from 10% to 70% w/w of the composition.

More preferably the composition in accordance with the present invention comprises nimesulide as an active drug from 40% to 60% w/w of the composition, one or more sustaining materials from 8% to 20% w/w of the

composition and pharmaceutical excipients from 30% to 60% w/w of the composition.

In a preferred embodiment of the invention the composition consists of bilayer tablets wherein the active agent may be present in one or both layers. The bilayer tablets may be coated or uncoated. The coating may be semi-permeable type membrane. Further, the semi-permeable coat may have an orifice drilled through it on the drug layer side to provide passage for constant release of drug.

In another aspect of the invention the coating may be of microporous type through which the drug release takes place at constant rate.

In another aspect of the invention the bilayer tablet dosage form may have a first layer which gives fast release of the drug, and a second layer which gives extended release of the drug.

The first fast release layer comprises materials like disintegrants, fillers, rapidly soluble/dispersible excipients and wetting agents. The second extended release layer comprises sustaining polymers binders wetting agents and fillers.

In another preferred embodiment of the invention, controlled release of Nimesulide from an oral formulation may be achieved by designing a gastroretentive system. It has been found by our group that Nimesulide is

absorbed mainly from the stomach and proximal part of small intestine. It is not well absorbed from distal part of small intestine and large intestine, leading to loss of bioavailability. Drugs having such characteristics require gastroretentive systems wherein the residence time of the drug is increased in the stomach, duodenum, jejunum or ileum. Gastroretentive characteristics may be incorporated into the Dosage Form/Drug Delivery system by techniques such as :

- a) Treating the drug with polymers having specific affinity to bind with gastric mucosa.
- b) Reduced specific gravity and floatation
- c) Particle Size
- d) Chemicals which delay gastric emptying and the like.

The compositions may be in the form of tablets, effervescent tablets, dispersible tablets, hard shell capsules, soft shell capsules, dry powder filled in sachet, dry powder for reconstitution or suspension.

The sustaining polymers preferably are hydrophilic in nature and present in a blend of fast and slow hydrating polymers.

The sustaining materials comprise materials which are non-toxic and pharmaceutically acceptable. These may be natural, semi-synthetic, synthetic or man-modified. Suitable materials include cellulose and cellulose derivatives like microcrystalline cellulose, methyl cellulose, ethyl cellulose, hydroxypropylmethyl cellulose, hydroxypropyl cellulose, cellulose acetate

WO 01/22791

PCT/IN 00/00094

phthalate, cellulose acetate, cellulose acetate butyrate, cellulose acetate propionate, cellulose acetate trimellitate, cellulose carboxymethyl ethers and their salts, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate.

Polyethylene; Polyquaternium-1; Polyvinyl acetate (homopolymer); Polyvinyl acetate phthalate; Propylene glycol alginate; PVM/MA copolymer; PVP/dimethiconylacrylate/polycarbamyl/polyglycolester; PVP/dimethylaminoethylmethacrylate copolymer; PVP/dimethylaminoethylmethacrylate/polycarbamyl polyglycol ester; PVP/polycarbamyl polyglycol ester; PVP/VA copolymer

Lanolin and lanolin derivatives, glyceryl monostearate, stearic acid, paraffins, beeswax, carnauba wax, Tribehenin.

Polyalkylene polyols like polyethylene glycols.

Gelatin and gelatin derivatives.

Alginates. Carbomers. Polycarbophils.

Methacrylic acid copolymers.

WO 01/22791

PCT/IN 00/00094

Carrageenans, pectins, chitosans, cyclodextrins, lecithins.

Natural and synthetic gums containing galactomannans like xanthan gum, tragacanth, acacia, agar, guar gum, etc.

Pharmaceutical excipients as used in the composition are selected from the group of excipients generally used by persons skilled in the art e.g. fillers, bulking agent, colourants, stabilizers, preservatives, lubricants, glidants, chelating agents and the like.

Preferably the composition also comprises release modifiers. Such release modifiers are selected from the groups wetting agents, solubilizers, surfactants, plasticizers, solvents, pore formers, pH modifiers and tonicity adjusting agents.

Suitable example of such ingredients include :

Reaction products of natural and hydrogenated vegetable oils and ethylene glycol e.g. polyoxyethylene glycolated natural or hydrogenated castor oil such as those available under the trade name Cremophor.

Other suitable products include polyoxyethylene sorbitan fatty acid esters e.g. of the type available under the trade name TWEEN.

Polyoxyethylene fatty acid esters e.g. MYRJ and CETIOL HE.

Polyoxyethylene polyoxypropylene copolymers e.g. PLURONIC and

Polyoxyethylene polyoxypropylene block copolymers e.g. POLOXAMER.

Diocylsodiumsulfosuccinate, sodium lauryl sulphate.

Propylene glycol mono- and di- fatty acid esters e.g. MIGLYOL 840.

Bile salts e.g alkali metals salts e.g. sodium taurocholate.

Polyethylene glycols, propylene glycol, triacetin, diacetin, diethyl phthalate, dibutyl phthalate, castor oil, triethyl citrate dibutyl sebacate.

Sodium chloride, potassium chloride, lactose, mannitol, sucrose, sorbitol.

Sodium hydroxide, potassium hydroxide, sodium bicarbonate, sodium citrate, citric acid, hydrochloric acid, lactic acid, tartaric acid, malic acid.

The calculation of dose of nimesulide for once-a-day controlled release dosage form was done on the basis of its pharmacokinetic parameters using the following equation :

$$\text{Dose} = C_p \times V_d \times K_{el} \times T$$

$C_p$  = Effective plasma concentration, 3.0 mg/L

$V_d$  = Apparent Volume of distribution, 15.6 L

$K_{el}$  = Elimination Rate constant,  $0.166 \text{ h}^{-1}$

$T$  = Desired Duration of action, 24 hrs

Based on the above equation the dose was calculated to be 207.0 mg.

The compositions of the present invention have another added advantage that once – a – day dosage form of Nimesulide may be combined with another suitable long – acting drug to have synergistic activity. The other



drug may be present in non-controlled release form. Such drugs may be selected from following categories :

- (i) Antihistaminics e.g. Cetirizine Dihydrochloride.
- (ii) Antispasmodics e.g. Pitofenone Hydrochloride, Hyoscine Hydrobromide.
- (iii) Antiasthmatics e.g. Ketotifen, Salbutamol.

The foregoing examples are illustrative embodiments of the invention and are merely exemplary. A person skilled in the art may make variations and modifications without depending from the spirit and scope of the invention. All such modifications and variation are intended to be included within the scope of the invention as discuss in this specifications.

**Example 1** Controlled release matrix tablet type

(i) Nimesulide (micronized)	-	200 mg
(ii) Lactose	-	73 mg
(iii) Hydroxypropylmethyl Cellulose	-	70 mg
(iv) Magnesium Stearate	-	3.5 mg
(v) Purified Talc	-	3.5 mg

Blend (i), (ii), (iii), (iv) and (v) after sifting through mesh no. 30 (BSS).  
Compress into tablets.

The results of Dissolution Release Profile of Nimesulide CR Tablets based on example 1 are given below :

**Table 1**

Time	Mean	SD
30 mins.	4.2	$\pm 1.36$
1 hr	7.9	$\pm 1.02$
2 hrs	16.4	$\pm 1.74$
3 hrs	25.8	$\pm 1.28$
4 hrs	34.2	$\pm 1.71$
6 hrs	50.8	$\pm 2.44$
8 hrs	65.9	$\pm 1.86$
10 hrs	74.9	$\pm 0.97$
12 hrs	85.8	$\pm 2.34$
14 hrs	93.5	$\pm 2.49$
16 hrs	96.7	$\pm 2.16$
18 hrs	97.1	$\pm 1.08$
19 hrs	98.8	$\pm 1.32$

The dissolution profile as given in table 1 of the nimesulide sustained release tablet should not be construed to limit the scope of the invention. Variations to the dissolution profile can be possible depending upon the dosage requirements without departing from the spirit of the invention.

**Example 2** Extended release membrane diffusion controlled tablet type

(i) Nimesulide (micronized)	-	200 mg
(ii) Microcrystalline Cellulose	-	60 mg
(iii) Lactose	-	60 mg
(iv) Maize Starch	-	10 mg
(v) Purified Talc	-	3.5 mg
(vi) Ethyl Cellulose	-	10 mg
(As Aqueous Dispersion)		
(vii) Polyethylene Glycol	-	3.5 mg

WO 01/22791

PCT/IN 00/00094

Blend (i), (ii) and (iii) and granulate with starch paste and dry the granules. Sift through mesh no. 22 (BSS). Lubricate with Talc. Compress into tablets. Coat the tablets with Ethyl Cellulose using Polyethylene Glycol as a channel former.

**Example 3** Sustained release bead type

(i) Non – Pareil Beads	-	347 mg
(ii) Nimesulide	-	200 mg
(iii) Mannitol	-	30 mg
(iv) Lactose	-	30 mg
(v) Polyvinyl Pyrrolidone	-	20 mg
(vi) Purified Talc	-	15 mg
(vii) Ethyl Cellulose	-	7 mg
(viii) Diethyl Phthalate	-	1.4 mg

Coat the non-pareil beads with blend of (ii), (iii) and (iv) using (v) as a binder in a conventional or fluidized bed coater. Talc may be dusted onto the beads. Final coating is given with Ethyl Cellulose using (viii) as plasticizer.

**Example 4** Osmotically controlled constant release type device

Upper Layer

(i) Nimesulide (micronized)	-	200 mg
(ii) Sodium Hydroxide	-	15 mg
(iii) Lactose	-	34 mg
(iv) Sodium Chloride	-	30 mg

WO 01/22791

PCT/IN 00/00094

(v) Polyvinyl Pyrrolidone	-	6 mg
(vi) Polyethylene Oxide	-	1.5 mg
<u>Lower Layer</u>		
(vii) Polyethylene Oxide	-	22 mg
(viii) Hydroxypropylmethyl Cellulose	-	1.8 mg
(ix) Sodium Chloride	-	20 mg
(x) Dichloromethane	-	q.s (Lost in processing)
<u>Semi-permeable Coat</u>		
(xi) Cellulose Acetate	-	30 mg
(xii) Triacetin	-	1 mg
(xiii) Acetone	-	q.s (Lost in processing)
(xiv) Water	-	q.s (Lost in processing)

Blend finely powdered (i), (ii), (iii), (iv) and (vi). Granulate with aqueous solution of (v). Granulate the blend of (vii) and (ix) with dispersion of (viii) in (x). Compress the two granulates into bilayer tablets and coat with the dispersion of (xii) and (xiii) in aqueous acetone. Finally, drill a hole in the drug layer (Upper layer) through which the drug is released in a controlled fashion due to osmotic pressure.

The results of Dissolution Release Profile of Nimesulide CR Tablets based on example 4 are given below :

**Table 2**

Time	Mean	SD
2 hours	5.16	± 0.53
4 hours	16.75	± 1.68
6 hours	34.90	± 2.26
8 hours	45.75	± 2.26

WO 01/22791

PCT/IN 00/00094

10 hours	56.00	$\pm 4.36$
12 hours	67.85	$\pm 4.40$
14 hours	79.16	$\pm 5.03$
14 hours	90.25	$\pm 3.68$
18 hours	101.16	$\pm 3.53$

**Example 5** Coated capsule type

(i) Nimesulide (micronized)	-	200 mg
(ii) Microcrystalline Cellulose	-	88.4 mg
(iii) Lactose		70 mg
(iv) Polyvinyl Pyrrolidone	-	7 mg
(v) Magnesium Stearate	-	3.9 mg
(vi) Ethyl Cellulose	-	20 mg
(vii) Polyethylene Glycol	-	0.7 mg
(viii) Alcohol : Dichloromethane (1 : 2)	-	q.s (Lost in processing)
(ix) Empty Gelatin Capsule (Size '1')		

Blend (i), (ii), (iii), (iv) and (v) and fill into empty gelatin capsule size '1'. Coat the capsules with dispersion of (vi) and (vii) in (viii).

**Example 6** pH dependent delayed release type

(i) Nimesulide (micronized)	-	100 mg
(ii) Microcrystalline Cellulose	-	150 mg
(iii) Lactose	-	76 mg
(iv) Polyoxyl 40 Hydrogenated Castor Oil	-	7 mg
(v) Polyvinyl Pyrrolidone	-	10 mg
(vi) Magnesium Stearate	-	3.5 mg

WO 01/22791

PCT/IN 00/00094

(vii) Purified Talc	-	3.5 mg
(viii) Cellulose Acetate Phthalate	-	28 mg
(ix) Diethyl Phthalate	-	2 mg
(x) Water	-	q.s (Lost in processing)
(xi) Alcohol : Dichloromethane (1:2)	-	q.s (Lost in processing)

Granulate the blend of (i), (ii) and (iii) with solution of (iv) and (v) in water.

Blend the granules with (vi) and (vii). Compress into tablets. Coat with the dispersion of (viii) and (ix) in (xi).

#### Example 7 Timed release bead type

(i) Nimesulide (micronized)	100 mg	100 mg	100 mg
(ii) Microcrystalline Cellulose	200 mg	200 mg	200 mg
(iii) Lactose	50 mg	42 mg	35mg
(iv) Polyvinyl Pyrrolidone	10 mg	10 mg	10 mg
(v) Water	q.s	q.s	q.s
(vi) Ammonio Methacrylate			
Copolymer Type B	10 mg	18 mg	25 mg
(Eudragit RS)			
(vii) Diacetin	0.5 mg	0.5 mg	0.5 mg
(viii) Water : Acetone (1:9)	q.s	q.s	q.s

Procedure :

In this composition 3 types of beads are prepared which are coated with different amounts of (vi) to give a timed profile of the drug. Beads are prepared by blending and spheronizing (I), (ii) and (iii) using aqueous solution of (iv). The dried beads are coated with dispersion of (vi) and (vii) in (viii). The 3 different beads are blended together in a fixed ratio to obtain the required release profile.

#### **Example 8 Nimesulide CR + Cetirizine Bilayered Tablets**

##### Nimesulide Layer

(i) Nimesulide (micronized)	-	200 mg
(ii) Lactose	-	106.5 mg
(iii) Polyoxyl 40 Hydrogenated Castor Oil	-	2.0 mg
(iv) Hydroxypropylmethylcellulose	-	31.5 mg
(v) Magnesium Stearate	-	2.0 mg
(vi) Colloidal Silicon Dioxide	-	2.0 mg

##### Cetirizine Layer

(vi) Colloidal Silicon Dioxide	-	2.0 mg
(vii) Cetirizine Dihydrochloride	-	10.0 mg
(viii) Lactose	-	105.0 mg
(ix) Microcrystalline Cellulose	-	25.0 mg
(x) Starch	-	5.0 mg
(xi) Croscarmellose Sodium	-	3.0 mg
(xii) Magnesium Stearate	-	2.0 mg

Blend the components of the two layers separately and compress into bilayer tablets.

**Example 9 Osmotically controlled constant release system**Active Layer

(i)	Nimesulide (micronized)	-	200.0 mg
(ii)	Polyethylene oxide	-	116.5 mg
(iii)	Hydroxypropylmethyl cellulose	-	10.0 mg
(iv)	Sodium chloride	-	10.0 mg
(v)	Magnesium stearate	-	2.5 mg

Push layer

(vi)	Polyethylene oxide	-	140.0 mg
(vii)	Sodium chloride	-	50.0 mg
(viii)	Hydroxypropylmethyl cellulose	-	9.5 mg
(ix)	Magnesium stearate	-	0.5 mg
(x)	Iron oxide red	-	1.0 mg

Functional coating

(xi)	Cellulose acetate	-	45.0 mg
(xii)	Polyethylene glycol	-	5.0 mg
(xiii)	Acetone	-	Lost in processing

Non-functional coating

(xiv)	Titanium dioxide	-	2.0 mg
(xv)	Hydroxypropylmethyl cellulose	-	6.0 mg
(xvi)	Purified Talc	-	2.0 mg
(xvii)	Polyethylene glycol – 400	-	2.0 mg
(xviii)	Isopropyl Alcohol	-	Lost in processing
(xix)	Dichloromethane	-	Lost in processing



Procedure : Blend (I), (ii), (iii), (iv) and (v) in a double cone blender. Separately blend (vi), (vii), (viii) (ix) and (x). Compress into bilayer tablet using a suitable compression machine. Coat the tablets with the dispersion of (xi) and (xii) in (xiii). The tablets are further coated with the dispersion of (xiv), (xv), (xvi), (xvii) in mixture of (xviii) and (xix).

**Example 10 : Bilayer tablets having one fast release layer and one extended release layer**

Fast Release layer

(i)	Nimesulide (micronized)	-	100.0 mg
(ii)	Lactose	-	151.5 mg
(iii)	Starch	-	37.6 mg
(iv)	Colloidal silicon Dioxide	-	11.0 mg
(v)	Povidone K-30	-	8.5 mg
(vi)	Docusate Sodium	-	6.8 mg
(vii)	Polysorbate 80	-	1.0 g
(viii)	Magnesium Stearate	-	1.6 mg
(ix)	Croscarmellose Sodium	-	22.0 mg
(x)	Water	-	Lost in processing

Extended Release Layer

(xi)	Nimesulide (micronized)	-	100.0 mg
(xii)	Lactose	-	200.0 mg
(xiii)	Hydroxypropylmethyl cellulose K100LV	-	23.0 mg
(xiv)	Hydroxypropylmethyl cellulose K4MCR	-	100.0 mg
(xv)	Povidone K-30	-	9.0 mg

WO 01/22791

PCT/IN 00/00094

(xvi)	Docusate Sodium	-	4.5 mg
(xvii)	Magnesium Stearate	-	4.5 mg
(xviii)	Colloidal Silicon Dioxide	-	4.5 mg
(xix)	Sodium Lauryl Sulphate	-	4.5 mg
(xx)	Isopropyl Alcohol	-	Lost in processing

**Procedure :**

Blend 1 : Blend (I), (ii), (iii) and (iv) and granulate with solution of (v) and (vi) in (x). Dry the granules and blend with (viii) and (ix).

Blend 2 : Blend (xi), (xii), (xiii) and (xiv) and granulate with solution of ((xv) and (xvi) in (xx). Dry the granules and mix with (xvii), (xviii) and (xix).

Compress into bilayer tablets using a suitable compression machine.

**Example 11 : Bilayer tablets having one fast release layer containing drug in complexed form and one extended release layer**

**A. Fast Release layer**

(i)	Nimesulide (micronized)	-	100.0 mg
(ii)	β-cyclodextrin	-	400.0 mg
(iii)	Starch	-	70.0 mg
(iv)	Povidone K-30	-	7.5 mg
(v)	Croscarmellose Sodium	-	20.0 mg
(vi)	Magnesium Stearate	-	2.5 mg

**B. Extended Release Layer**

(vii)	Nimesulide (micronized)	-	100.0 mg
-------	-------------------------	---	----------

WO 01/22791

PCT/IN 00/00094

(viii)	Lactose	-	200.0 mg
(ix)	Hydroxypropylmethyl cellulose K100LV	-	23.0 mg
(x)	Hydroxypropylmethyl cellulose K4MCR	-	100.0 mg
(xi)	Povidone K-30	-	9.0 mg
(xii)	Magnesium Stearate	-	4.5 mg
(xiii)	Colloidal Silicon Dioxide	-	4.5 mg
(xiv)	Docusate Sodium	-	4.5 mg

#### Procedure :

##### Layer – I

1. Mix (i) and (ii), co-mill under specific conditions favouring complexation using ball mill to prepare a complex.
2. Mix complex of step 1 with (iii) and granulate with a solution of (iv) in water.
3. Dry the granules at 40° - 50°C.
4. Size the granules & mix with (v) and (vi)

##### Layer – II

1. Mix (vii), (viii), (ix) and (x). Granulate with a solution of (xi) and (xiv).
2. Dry the granules at 40° - 50°C.
3. Size the granules & mix with (xii) and (xiii).
4. Compress the two layers into bilayered tablets using suitable compression machine.

#### Example 12 Controlled release mucoadhesive granules

WO 01/22791

PCT/IN 00/00094

i)	Nimesulide	-	200 mg
ii)	Hydrogenated vegetable oil	-	45 mg
iii)	Carbopol 974P	-	50 mg
iv)	Dibutyl sebacate	-	10 mg

1. Melt (ii) at 60°C and granulate (i) with it. Pass the granules through mesh no. 60 BSS.
2. Disperse (iii) and (iv) in isopropyl alcohol.
3. Coat granules of step (1) with the solution of step (2) in fluidized bed coater.

The resultant granules may incorporated into tablets, may be filled in hard gelatin/soft gelatin capsules or may be prepared into dry syrups for reconstitution.

### Example 13 Gastroretentive granules

i)	Nimesulide	-	200 mg
ii)	Hydroxypropylmethyl cellulose	-	20 mg
iii)	Ethyl cellulose	-	50 mg
iv)	Triethyl citrate	-	12.5 mg
v)	Carbopol 934P	-	50 mg

1. Granulate (i) with an aqueous dispersion of (ii). Dry the granules and sift through mesh no. 60 BSS.

WO 01/22791

PCT/IN 00/00094

2. Dissolve (iii) in isopropyl alcohol/dichloromethane and add (iv).
3. Coat the granules of step (1) with solution of step (2) in Fluid bed coater.
4. Apply another coat of Carbopol 934P onto the coated granules of step (3) in fluidized bed coater.

The resultant granules may incorporated into tablets, may be filled in hard gelatin/soft gelatin capsules or may be prepared into dry syrups for reconstitution.

**Example 14 Mucoadhesive granules**

i)	Nimesulide	-	200 mg
ii)	Chitosan	-	40 mg
iii)	Polyethylene glycol	-	40 mg
iv)	Carbopol 934P	-	50 mg

1. Granulate (i) with aqueous dispersion of (ii) using (iii) as the plastisizer. Pass the granules through mesh no. 60 BSS.
2. Disperse (iv) in Isopropyl Alcohol/Water.
3. Coat the granules with (iv) on fluidized bed coater.

The resultant granules may incorporated into tablets, may be filled in hard gelatin/soft gelatin capsules or may be prepared into dry syrups for reconstitution.

**Example 15 Controlled release suspension (Dry powder for reconstitution)**

i)	Nimesulide Granules as per example		
	12,13 or 14 equivalent to Nimesulide	-	200 mg
ii)	Sucrose	-	500 mg
iii)	Xanthan Gum	-	10 mg
iv)	Flavour orange	-	5 mg
v)	Citric acid	-	12.5 mg
vi)	Sodium citrate	-	5 mg

1. Prepare a dry powder for reconstitution by mixing (i) with (ii), (iii), (iv), (v) and (vi).
2. The powder may be reconstituted with water to contain 25 mg Nimesulide per ml.

**Example 16 Mucoadhesive Tablets**

i)	Nimesulide granules as per examples		
	12, 13 or 14 equivalent to Nimesulide	-	200 mg
ii)	Microcrystalline Cellulose	-	250 mg
iii)	Sodium Lauryl Sulphate	-	1 mg
iv)	Polyvinyl Pyrrolidone	-	15 mg
v)	Magnesium Stearate	-	6 mg

WO 01/22791

PCT/IN 00/0094

vi)	Purified Talc	-	6 mg
-----	---------------	---	------

Mix (i) with (ii), (iii), (iv), (v) and (vi) after sifting through mesh no. 30 (BSS).

Compress into tablets.

#### Example 17 Mucoadhesive Effervescent Tablets

i)	Nimesulide granules given in examples		
	12, 13 or 14 equivalent to Nimesulide	-	200 mg
ii)	Sodium Bicarbonate	-	400 mg
iii)	Polyethylene Glycol 6000	-	20 mg
iv)	Magnesium Stearate	-	10 mg
v)	Purified Talc	-	10 mg

1. Mix (I) with (ii), (iii), (iv) and (v) after sifting through mesh no. 30 (BSS).

2. Compress into tablets.

#### Example 18 Gastroretentive Mucoadhesive Beads

i)	Non – Pareil Beads	-	350 mg
ii)	Nimesulide	-	200 mg
iii)	Lactose	-	30 mg
iv)	Ethyl Cellulose	-	350 mg
v)	Diethyl Phthalate	-	5 mg
vi)	Carbopol 934	-	50 mg

1. Coat the non-poreil beads with (ii), (iii) and (iv) using (v) as the plasticizer.
2. Disperse (vi) in Isopropyl Alcohol/Water and coat the drug loaded beads of Step – 1 with above solution in Fluidized Bed Coater.

**Example 19 Gastric mucoadhesive Nimesulide controlled release microspheres**

i)	Nimesulide	-	200 mg
ii)	Chitosan (Mol. Wt. 2,000,000)	-	40 mg
iii)	Sorbitan monooleate	-	0.5 mg
iv)	Liquid Paraffin (light)	-	25 ml
v)	Acetic acid	-	1 ml
vi)	Purified water	-	15 ml
vii)	Lectin (wheat germ agglutinin)	-	2.25 mg
viii)	Glutaraldehyde	-	0.1 ml
ix)	Toluene	-	2 ml

1. Dissolve (ii) in solution of (v) & (vi)
2. Disperse (i) in solution of (iii) & (iv)
3. Prepare an emulsion of solution of step-1 in dispersion of step-2 by continuous stirring.
4. Add solution of (viii) in (ix) to the emulsion of step-3. Continue stirring for 3 hours to prepare microspheres.



5. Filter the microspheres of step-4, wash and dry.
6. Resuspend microspheres of step-5 in pH 7.4 phosphate buffer and add (vii). Incubate overnight.
7. Centrifuge to separate microsphere – lectin conjugates from free lectins.

**Example 20 gastric mucoadhesive Bilayered tablets**

i)	Nimesulide (micronized)	-	200mg
ii)	Hydroxypropylmethyl Cellulose	-	40mg
iii)	Hydroxyethyl cellulose	-	10mg
iv)	Lactose	-	100mg
v)	Povidone	-	5mg
vi)	Magnesium Stearate	-	5mg
vii)	Polycarbophil	-	75mg

( Trade Name : Noveon AA1)

viii)	Lactose	-	150mg
ix)	Hydroxypropylmethyl cellulose	-	20mg
x)	Magnesium stearate	-	2.5mg
xi)	Isopropyl alcohol	-	q.s. lost in processing
xii)	Dichloromethane	-	q.s. lost in processing

1. Sift (i), (ii), (iii) and (iv) through sieve of mesh no. 40 and mix.
2. Dissolve (v) in (xi).
3. Granulate blend of step 1 with solution of step 2 and dry the granules.
4. Pass the granules through sieve of mesh no. 20.

5. Lubricate the granules of step 4 with (vi).
6. Separately sift (vii) and (viii) through sieve of mesh no. 40 and mix.
7. Dissolve (ix) in a mixture (xi) and (xii).
8. Granulate the blend of step 6 with solution of step 7.
9. Dry the granules and pass through sieve of mesh no. 20.
10. Lubricate the granules of step 9 with (x).
11. Compress the granules of step 5 and step 10 into bilayered tablets.

**Example 21 Hydrodynamically Balanced System**

i)	Nimesulide (micronized)	-	200mg
ii)	Carbomer 934P	-	25mg
(Trade name : Carbopol 934P)			
iii)	Hydroxypropyl Cellulose	-	50mg
iv)	Xanthan Gum	-	20mg
v)	Sodium Aliginate	-	15mg
vi)	Sodium Bicarbonate	-	100mg
vii)	Lactose	-	100mg
viii)	Povidone	-	10mg
ix)	Magnesium Stearate	-	5mg
x)	Collordal Silicon Dioxide	-	2.5mg
xi)	Talc	-	2.5 mg
xii)	Isopropyl alcohol	-	q.s. lost in processing

1. Sift (i), (ii), (iii), (iv), (v), (vi) and (vii) through sieve of mesh no. 40 and mix.

2. Dissolve (viii) in (xii) and granulate blend of step 1.
3. Dry the granules and pass through sieve of mesh no. 20.
4. Lubricate with (ix), (x) and (xi) and compress into tablets.

#### **Example 22 Intestinal Targeting of Mucoadhesive Tablets**

1.	Nimesulide ( micronized)	-	200mg
ii)	Lactose	-	100mg
iii)	Hydroxypropyl methylcellulose	-	50mg
iv)	Polycarbophil	-	50mg
	( Trade name : Noveon AA1)		
v)	Povidone/ Vinyl acetate copolymer	-	10mg
vi)	Magnesium Stearate	-	5mg
vii)	Talc	-	5mg
viii)	Methacrylic Acid Copolymer	-	33mg
	(Trade name : Eudragit L 100)		
ix)	Triethyl Citrate	-	7mg
x)	Talc	-	5mg
xi)	Colour Lake of Quinoline Yellow	-	1.5mg
xii)	Isopropyl alcohol	-	q.s. Lost in processing
xiii)	Dichloromethane	-	q.s. lost in processing

#### **Core Tablets**

1. Sift (i),(ii), (iii) and (iv) through sieve of mesh no. 40 and mix.
2. Dissolve (v) in (xii) and granulate blend of step1.
3. Dry the granules and pass through sieve of mesh no. 20.
4. Lebricate with (vi), (vii) and compress into tablets.

#### **Enteric Coating**

1. Dissolve (viii) and (ix) in a mixture of (xii) and (xiii)

2. Disperse (x) and (xi) in the solution of step 1.
3. Coat the tablets with the solution of step 2 till enteric effect is obtained.

**Example 23 Controlled Release Tablets**

i)	Nimesulide(micronized)	-	200mg
ii)	Lactose	-	100mg
iii)	Polyethylene oxide	-	50mg
iv)	Polyvinyl alcohol (mol. Wt. 1,400,00)	-	25 mg
v)	Polyvinyl acetate	-	20mg
vi)	Isopropyl alcohol	-	q.s. lost in processing
vii)	Magnesium stearate	-	2.5mg
viii)	Colloidal silicon dioxide	-	2.5 mg

1. Sift (i), (ii), (iii) and (iv) through sieve of mesh no. 30 (BSS) and blend.
2. Dissolve (v) in (vi) and granulate the blend of step -1.
3. dry the granules and pass through sieve of mesh no. 20 (BSS).
4. Lubricate with (vii) and (viii) and compress into tablets.

## We claim :

1. A controlled release pharmaceutical composition of Nimesulide which comprises nimesulide as an active drug upto 99% w/w of the composition, one or more release controlling materials from 0.1% to 99% w/w of the composition and pharmaceutical excipients from 0% to 90% w/w of the composition.
2. A controlled release pharmaceutical composition of nimesulide as claimed in claim 1 which comprises nimesulide as an active drug from 20% to 70% w/w of the composition, one or more sustaining materials from 5% to 65% w/w of the composition and pharmaceutical excipients from 10% to 70% w/w of the composition.
3. A controlled release pharmaceutical composition of nimesulide as claimed in claim 1 and 2 wherein there is no loss of bioavailability in comparison to an immediate release composition.
4. A controlled release pharmaceutical composition of nimesulide as claimed in claim 1 to 3 wherein the sustaining materials are selected from the group comprising cellulose and cellulose derivatives, waxes, carbomers, polyalkylene polyols, polycarbophils, methacrylic acid derivatives, gelatins, gums, polyethylene oxides.
5. The composition as claimed in claim 1 which further comprises release modifiers selected from the group comprising wetting agents, solubilizers, surfactants, plasticizers, pore formers, pH modifiers and tonicity adjusting agents.
6. A controlled release pharmaceutical composition as claimed in claim 1 which is a gastroretentive system wherein the residence time of the drug is increased in the stomach, duodenum, jejunum or ileum.
7. A composition as claimed in claim 6 wherein gastroretention of Nimesulide is achieved by mucoadhesion, floatation and/or reducing gastrointestinal motility.

8. A composition as claimed in claim 7 wherein mucoadhesion is achieved by treating Nimesulide with polymers having affinity for gastrointestinal mucosa comprising Polycarbophils, carbomers, alginates, Cellulose and Cellulose derivatives, Chitosan, Gums and Lectins.
9. A composition as claimed in claim 7 wherein floatation is achieved by adding to the composition gas-generating materials comprising sodium bicarbonate, sodium carbonate, calcium carbonate and potassium carbonate alone or in combination with acidic substances comprising hydrochloric acid, citric acid, fumeric acid, malic acid, maleic acid, ascorbic acid and tartaric acid.
10. A composition as claimed in claim 7 wherein gastrointestinal motility is reduced by materials comprising fats, fatty acids and transesterification products of fats and fatty acids with polyols.
11. A process for the manufacture of controlled release compositions of Nimesulide which comprises mixing together under conventional conditions of temperature and pressure – nimesulide as an active drug upto 99% w/w of the composition, one or more release controlling materials from 0.1% to 99% w/w of the composition and pharmaceutical excipients 0% to 90% w/w of the composition
12. A controlled release pharmaceutical composition of nimesulide substantially as herein described with reference to foregoing description and the accompanying examples
13. A process for the manufacture of controlled release compositions of nimesulide substantially as herein described with reference to foregoing description and accompanying examples.

## ABSTRACT

A Controlled release pharmaceutical composition of Nimesulide comprising Nimesulide, as an active drug, one or more, sustaining materials and pharmaceutical excipients formulated into a controlled release once-a-day oral dosage form.